Chapter 44

Appendix List of condition assessments

List of condition assessments (ranked as very good, good, poor, very poor) for estuaries and deltas in different regions, and the trend in condition over time intervals specified). Year of assessment, the number of estuaries or deltas included and references are indicated for each region. Indicative grading statements used to define the condition are as follows:

Very Good = Estuarine habitats are essentially structurally and functionally intact and able to support all dependent species. Only a few, if any, species populations have declined as a result of human activities or declining environmental conditions. There are no significant changes in physical-chemical

					coastal habitat index, and a fish tissue contaminants index.
United States, Hudson River Estuary	1	Very Poor (2012)	Stable (2005-2010)	New York and New Jersey Harbor and Estuary Program (2012)	Assessment of marine habitat, fish, birds and pollution impacts.
United States, SE Coast	2	Good (2000-02)			

Caribbean river basins	>4	Poor (2002)	Declining	Kjerfve et al. (2002)	Integrated LOICZ assessment coverin Caroni River (Trinidad), Kingston Harbour (Jamaica), Parque Nacional Morrocoy (Venezuela) and Magdelena River (Costa Rica)
Uruguay, Río de la Plata estuarine environment Montevideo Bay (including Montevideo Harbour)	1	Poor to very poor (1997–2010)	Improving	Muniz et al., (2002, 2004, 2005, 2006, 2011, 2012); García-Rodríguez et al. (2010) Burone et al. (2006, 2011) Danulat et al (2002) Gómez-Erache et al. (2001)	Biotic indices based on benthic communities; coastal eutrophication; heavy metals and PAH in sediments
Uruguay, Río de la Plata estuarine environment (seaward), Montevideo Coastal zone	1	Poor to good (1997–2012)	Declining	Muniz et al., (2002, 2004, y	

(coastal lagoons and sub-estuaries)	Conde & Rodríguez-Gallego (2002)	2

Russian Federation, North Dvina River Estuary	1	Very Poor (1990-2006)	Declining (2006)	Gordeev et al. (2006)	LOICZ-DPSIR approach based mainly o water quality (heavy metals and hydrocarbons, acidification and radionuclide contamination)
Russian Federation, Small rivers of the Kola					

rivers					
India, small western rivers flowing into the Arabian Sea	?	Very Poor (2005)	Declining	Ramesh et al. (2009)	LOICZ - DPSIR approach - integrated assessment
India and Pakistan, Indus River	1	Very Poor (2005)	Declining	Ramesh et al. (2009)	LOICZ - DPSIR approach - integrated assessment
India, Sabarmati River	1	Very Poor (2014)	Declining (2009-2014)	Haldar et al (2014)	Turbidity, dissolved oxygen, BOD, phenol and petroleum hydrocarbons, phytoplankton and total and selective bacterial count.
South China Sea estuaries and deltas		Poor (2012)	Declining (2005-2012)	Ward (2012)	Integrated assessment
China, Changjiang (Yangtze) River estuary	1	Very Poor (2003)	Declining (1999-2010)	Xiao et al. (2007); Wang (2007); Liu et al (2013)	Eutrophication - ASSETS and AMBI index methods
China, Jiaozhou Bay	1	Very Poor (2006)	Declining	Bay	

Arab Waterway		(2012)		(2006); Mohamed et al. (2012)	production,changes in water quality, and specific populations of rare and endangered species
Republic of Korea, Gwangyang Bay	1	Good (2012)	Stable (2010-2012)	Kim et al., (2008); Lee et al., (2010); KIOST (2013)	Organochlorine pesticides; Water quality; the carbon isotope ratio of particulate organic matter and sediment, and the nutrients limiting phytoplankton growth; marine ecosystem health index (MEHI) based on water quality, sediment quality, plankton, and benthos
Republic of Korea, Jinhae Bay	1	Poor (2012)	Stable (2010-2012)	Lim et al., (2012); KIOST (2013)	Sediment core records of C, N, $CaCO_3$, trace metals; MEHI
Japan, Mutsu Bay	1	Good (2012)	Declining (2004-12)	Environment Management Bureau, Ministry of the Environment (2005-13)	Environmental status evaluated by Dr. K. Kohata from Japanese Environmental Quality Standards (for Living Environment)
Japan, Toyama Bay	1	Very Good (2012)	Stable (2003-12)	Toyama Prefecture (2009, 2014)	Environmental status evaluated by Dr. K. Kohata from Japanese Environmental Quality Standards
Japan, Tokyo Bay					

Shimabara Bay				Environment (2013)	Quality Standards
Japan, Yatsushiro Sea	1	Good (2012)	Stable (2003-12)	Environment Management Bureau, Ministry of the Environment (2013)	Environmental status evaluated by Dr. K Kohata from Japanese Environmental Quality Standards

				Gu et al.(2013)	
Kenya, Tana an Athi- Sabaki estuaries	2	Good (2009)	Declining	UNEP/Nairobi Convention Secretariat and WIOMSA (2009)	Nutrient levels in the Tana and Athi-Sabaki estuaries are high; the threshold chemical contamination levels of both Tana and Athi-Sabaki rivers have not been

					mangrove and coral reef habitat, overfishing

		(2006)		Ezhova et al. (2012)	
Finland, Kvarken Archipelago Russian Federation, Neva Bay, Gulf of Finland	1	Good (2010)		Kotilainen et al., (2012)	Benthic habitat mapping

on the Skagerrak coas				areas/Coastalwaters/	
Norway, Nothern fjords		Good (2014)	Declining	www.environment.no /Topics/Marine areas/Coastalwaters/	
Antarctica					
Prydz Bay fjords	1	Very Good (2010)		O'Brien et al. (2012)	

Burone, L., Muniz, P. et al. (2011). Evolución paleoambiental de la Bahía de Montevideo (Uruguay) – bases para el establecimiento de un modelo ambiental. Libro: El Holoceno en la zona costera del Uruguay. pp: 197 - 227, Organizadores: F. García-Rodríguez Editorial: Tradinco , Montevideo. ISSN/ISBN: 9789974007574

Buruaem, L.M., de Castro, I.B., Hortellani, M.A., Taniguchi, S., Fillmann, G., Sasaki, S.T., Petti,

Institute of Zoology, Zoological Society of London, U.K. and WWF International, Gland, Switzerland, p. 49.

Defeo, O., Horta, S., Carranza, A., Lercari, D., de Álava, A., Gómez, J., Martínez, G., Lozoya, J.P., Celentano, E. (2009). Hacia un Manejo Ecosistémico de Pesquerías. Áreas Marinas Potegidas en Uruguay. Facultad de Ciencias, DINARA, Montevideo.

Department of Environment (2011). Australia State of the Environment 2011: Marine Environment. Canberra: Australian Department of Sustainability, Environment, Water, Population and Communities on behalf of the State of the Environment 2011 Committee. http://environment.gov.au/soe

Díaz-Jaramillo, M., Socowsky, R., Pardo, L.M., Monserrat, J.M., Barra, R. (2013). Biochemical responses and physiological status in the crab Hemigrapsus crenulatus (Crustacea, Varunidae) from high anthropogenically impacted estuary (Lenga, south-central Chile). Marine Environmental Research 83, 73-81.

Dupra, V., Smith, S.V., David, L.T., Waldron, H., Marshall Crossland, J.I., Crossland, C.J. (2002). Estuarine systems of Africa: carbon, nitrogen and phosphorus fluxes., LOICZ

Moscoso, J., Rudolph, A., Sepúlveda, R.D., Suárez, C. (2006). Effect of Temporary Closure of the Mouth of an Estuary on the Benthic Macroinfauna: Lenga, Chile, A Case Study. Bulletin of Environmental Contamination and Toxicology, 77, 484-491.

Muniz, P.; Venturini, N.; Martínez, A. (2002). Physico-chemical characteristics and pollutants of the benthic environment in the Montevideo coastal zone, Uruguay. Marine Pollution Bulletin, v.: 44, p.: 962 -968.

Muniz, P., Danulat, E.; Yannicelli, B. García-Alonso, J., Medina, G., Bícego, M. C., (2004). Assessment of contamination by heavy metals and petroleum hydrocarbons in sediments of Montevideo harbour (Uruguay). Environment International, v.: 29, p.: 1019