

Chapter 51 Biological Communities on Seamounts and Other Submarine Features Potentially Threatened by Disturbance

Contributors: J. Anthony Koslow, Peter Auster, Odd Aksel Bergstad, J. Murray Roberts, Alex Rogers, Michael Vecchione, Peter Harris, Jake Rice, Patricio Bernal (CoLead members)

1. Physical, chemical, and ecological characteristics

1.1 Seamounts

Seamounts are predominantly submerged volcanoes, mostly extinct, rising hundreds to thousands of metres above the surrounding seafloor. Some also arise through tectonic uplift. The conventional geological definition includes only features greater than 1000 m in height, with the term “knoll” often used to refer to features 100–1000 m in height (Yesson et al. 2011). However, seamounts and knolls do not appear to differ much ecologically, and human activity, such as fishing, occurs on both. We therefore include here all such features with heights > 100 m.

Only 6.5 per cent of the deep sea (2001) has been identified and guyot features > 1000 m in height and 138,412 (Harris et al. 2014) identified 10,234 seamount and guyot features under a stricter definition that restricted seamounts to conical forms. (Smith 1991; Wesselet al., 2010) At least half are in the Pacific, with fewer in the Atlantic, Indian, Southern, and Arctic Oceans. They cover approximately 4.7 per cent of the ocean floor, with an additional 16.6 per cent in total an area approximately the combined area of all continental shelves (Boyer et al. 2010; Yesson et al., 2011).

Local ocean circulation, amplifying and rectifying flows particularly near seamount summits, enhancing vertical mixing, and forming cells known as Taylor columns or cones. Some seamounts have many factors, including the size (height and diameter) of the

Where flows are sufficiently vigorous, they provide sufficient flow of organic matter to support suspension feeding organisms, such as corals and sponges. Such currents also winnow away the sediment, providing hard substrate necessary for most

such as Antarctic Intermediate Water and North Atlantic Deep Water (Koslow et al., 1994; Clark et al., 2010a and b). Whereas the dominant genera and families of deep demersal and midwater fishes tend to have global distribution, the dominant fish species on seamounts in different ocean basins are often from entirely different genera, families, and even orders. This indicates that seamount-associated fishes in different ocean basins were reproductively isolated and evolved independently. Their similar morphologies and adaptation to the seamount environment is a striking example of convergent evolution (Koslow, 1996).

Seamounts are the source of significant ecosystem services. In addition to their biodiversity, seamounts often host substantial aggregations of fishes, which have been subject to commercial fisheries. These include species for which seamounts are their primary environment as well as a larger number for which seamounts account for a smaller proportion of their global catch. Annual landings of primary seamount species

Ridges typically contain seamounts and sedimented slopes. Not surprisingly, similarities in the abundance, diversity, and species composition of ridge habitats are found

Monterey Bay Aquarium Research Institute (MBARI) Monterey Canyon have led to a renaissance in canyon studies (e.g. Huvenne and Davy, 2013). Whereas most canyons globally have received little or no scientific attention from any discipline, some individual canyons (e.g., Monterey – western North America, “The Gully” eastern North America, Kaikura New Zealand,

1.4 Trenches

Trenches are defined as “long, narrow, characteristically very deep and asymmetrical depressions of the seafloor, with relatively steep sides” (IHO

anemones and their mobile benthic associates (e.g., amphipods); communities near hydrothermal vents and cold seeps are dominated by metazoans dependent on

and the Chilean Rise. Catches were low nor were they sustainable (Clark et al. 2007).

In the Southern Ocean, seamounts were fished for nototheniids between 1974 and 1991. In the 1990s, the ridges, plateaus, and seamounts around remote Antarctic islands came to be heavily fished for Patagonian toothfish with trawls and longlines. Initially much illegal, unreported and unregulated (IUU) fishing occurred but has declined significantly since 1996 (Agnew et al. 2009).

Large scale industrial deepwater fisheries in the North Atlantic date to the development of redfish fisheries in the 1950s using both midwater and demersal trawls over the midAtlantic Ridge and on some plateaus. Redfish catches peaked at almost 400,000 tons in the 1950s and have declined considerably but several continue to support some harvest (Koslow et al. 2000; ICES 2013). Fisheries for roundnose grenadier and Greenland halibut first developed on the upper continental slopes of the Northwest Atlantic in the late 1960s, peaking at over 80,000 tons in 1971 and then rapidly declined and moved to the midAtlantic Ridge and Rockall Bank (1970s) (Clark et al. 2007).

Exploratory trawl fishing on seamounts in the Indian Ocean began in the 1970s targeting shallowwater redbait and rubyfish on the Southwest Indian Ocean Ridge, the Mozambique Ridge and the Madagascar Ridge (Rov, 2003; Clark et al 2007) and continued into the mid-1980s. In the late 1990s trawlers working on the Southwest Indian Ocean Ridge targeted deepwater species such as orange roughy, black cardinalfish, pelagic armorhead, oreosomatids and *Squalius* (Clark et al 2007) but the fishery rapidly collapsed (Gianni 2004). Fishing has shifted to the many ridges, seamounts and plateaus targeting a variety of species of deepfish and crustaceans (Clark et al. 2007; Bensch et al 2009; SWIOFC 2009).

Overall, deepwater demersal fisheries over the continental shelves, ridges, seamounts, and plateaus have landed between 800,000 and 1,000,000 t per annum from the mid-1960s to 1990s (Koslow et al, 2000) and annual landings on the order of 100,000 t since about 1990 (Clark et al 2007; Watson et al, 2007). The vast majority of seamount associated demersal fisheries have proven unsustainable, undergoing a boom-

- 4 Both the North Atlantic Fisheries Organization (NAFO) and North East Atlantic Fisheries Commission (NEAFC) in the North Atlantic set quotas for deep-sea stocks based on scientific assessments and have identified and closed to fishing areas that meet the Food and Agricultural Organization of the United Nations (FAO) criteria for vulnerable marine ecosystems
 - 4 The Southeast Atlantic Fisheries Organization (SEAFO) closed selected ridge sections and seamounts to fishing, restricted fisheries to certain subareas, and introduced catch quotas (TACs) for the fishes and deep-water crab targeted on seamounts.
 - 4 States which participated in the negotiations for the establishment of the North
- P108 708 Tm6.005 Tc Tc ,

concern, including mercury and many halogenated hydrocarbons (e.g., DDT, PCBs, and many other pesticides, herbicides, and industrial chemicals) are volatile and enter the ocean predominantly through the atmosphere. These are discussed in Chapter 20. As noted there, concentrations of persistent organic pollutants in deep-dwelling fish

loss and declining food availability. Midwater fishes, the primary food of many deepwater squid and fish species, including orange roughy, declined 60% during recent periods of low oxygen availability in the California Current (Koslow et al. 2011). Palaeoceanographic studies have pointed to the significance of perturbations in oxygen concentration in controlling deep coral occurrence in the Eastern Mediterranean (Fink

services are provided b

These characteristics lead to low productivity

5. Integrated assessment of the status of the habitat. Crossing and emergent conclusions

T

diversity and ecological connectivity. The synergistic influence of these factors is unknown at present.

Although it is heartening that some seamounts, ridges and other sensitive marine habitats are being protected by fishing closures, Marine Protected Areas and other actions, little scientific understanding of the efficacy of actions implemented to date and few studies to assess their impact exist. The connectivity between these habitats remains largely unknown, as are the factors that influence colonization, species succession, resilience and variability. Comparative studies of seamount, canyon, and continental margin habitats seem to indicate that many species are shared (but see Richer de Forges et al., 2000); however, community structure differs markedly and the factors influencing such differences remain unknown (McClain et al., 2009). Our starting point in attempting to understand a few of the factors influencing these differences is to assess the connectivity between these habitats and the factors that influence colonization, species succession, resilience and variability.

House, Moscow: 385 pp

- Bensch, A., Gianfranceschi, M., Gréboval, D., Sanders, J.S., Hjort, A. (2009). Worldwide review of bottom fisheries in the high seas. FAO Fisheries and Aquaculture Technical Paper 522. FAO, Rome: 145.
- Bergstad, O.A., Falkenheide, G., Astthorsson, O.S., Byrkjedal, I., Gebruk, A.V., Piatkowski, U., Priede, I.G., Santos, R.S., Vecchione, M., Lorange, P. (2008). Towards improved understanding of the diversity and abundance patterns of the mid-ocean ridge macro and megafauna. Deep Sea Research Part II: Topical Studies in Oceanography 55(1): 1-5.
- Blankenship, Williams, L.E. and Levin, L.A. (2009). Living Deep: a synopsis of hadal trench ecology. Marine Technology Society Journal 43(5): 137-143.
- Bograd, S.J., Castro, C.D., Lorenzo, E., Palacios, D.M., Bailey, H., Gilly, W., Chavez, F.P. (2008). Oxygen declines and the shoaling of the hypoxic boundary in the California Current. Geophysical Research Letters 35: L2607.
- Bruun, A.F. (1956). The abyssal fauna: its ecology, distribution and origin. Nature 177: 1105-1108.
- Castelin, M., Puillandre, N., Lozouet, P., Sysoev, A., de Forges, B., Samadi, S. (2011). Molluscan species richness and endemism on New Caledonian seamounts: Are they enhanced compared to adjacent slopes? Deep Sea Research Part I: Oceanographic Research Papers 58(6): 637-646.
- Clark, C.W. (1973). The economics of overexploitation. Science 181: 630-634.
- Clark, M.R. (2009). Deep-sea seamount fisheries: a review of global status and future prospects. Latin American Journal of Aquatic Research 37(3): 501-511.
- Clark, M.R., Althaus, F.,

Review of Marine Science 253278.

- Clark, M.R., Vinnichenko, I., Gordon, J.D.M., Beck, Bulat, G.Z., Kukharev, N.N., Kakora, A.F. (2007). Large scale distant water trawl fisheries on seamounts. In: T.J. Pitcher, J. Morato, T., Hart, P.J.B. et al (Eds). Seamounts: Ecology, Fisheries & Conservation Blackwell, Oxford: 363-399.
- Consalvey, M., Clark, M.R., Rowden, A.R., Stocks, K.I. (2010). Life on seamounts. In: A.D. McIntyre (Ed). Life in the world's oceans: diversity, distribution, and abundance Wiley-Blackwell, Blackwell Publishing Ltd, UK: 128.
- Dana, J.D. (1863) Manual of Geology Philadelphia: 798p.
- De Leo, F.C., Smith, R., Rowden, A.A., Bowden, D.A., Clark, M.R. (2010) Submarine canyons: hotspots of benthic biomass and productivity in the deep sea. Proceedings of the Royal Society B: Biological Sciences 277(1695): 2783-2792.
- Devillers, R., Presse, L., Grech, A., Kittinger, J.N., Edgar, G. J., Ward, T., Watson, R. (2014). Reinventing residual reserves in the sea: are we favouring ease of establishment over need for protection? Aquatic Conservation: Marine and Freshwater Ecosystems DOI: 10.1002/aqc.
- Devine, J.A., Baker, K., Eberhard, R.L. (2006). Deepsea fishes qualify as endangered Nature 439: 29.
- Etnoyer, P.J., Wood, S., Shirley, T.C. (2010). How large is the seamount biome? Oceanography 23: 206-209.
- FAO (2009) International Guidelines for the Management of Deep-sea Fisheries in the High Seas. Directives internationales sur la gestion de la pêche profonde en haute mer Directrices Internacionales para la Ordenación de las Pesquerías de Aguas Profundas en Alta Mar. FAO, Rome: 73 pp.
- Findlay, H.S., Hennige, J., Wicks, L.C., Navas, J.M., Woodward, E.M.S., Roberts, J.M. (2014). Fine scale nutrient and carbonate system dynamics around-water coral reefs in the northeast Atlantic. Nature Scientific Reports 4: 3671.
- Findlay, H.S., Wicks, Navas, J.M., Hennige, S., Huvenne, V., Woodward, E.M.S., Roberts, J.M. (2013). Tidal downwelling and implications for the carbon biogeochemistry of cold water corals in relation to future ocean acidification and warming. Global Change Biology 19: 2270-2289.
- Fink, H.G., Wienberg, Clebbeln, D., McGregor, H.V., Schmiedl, G., Taviani, M., Freiwald, A. (2012). Oxygen control on Holocene cold water coral development in the eastern Mediterranean Sea. Deep Sea Research Part I Oceanographic Research Papers 62s 8996.
- Form, A.U., Riebesell, (2012). Acclimation to ocean acidification during long term CO2 exposure in the cold water coral *Lophelia pertusa*. Global Change Biology 18: 843-853.

Fujii, T., Jamieson, A.Solan,M., Bagley,P.M.,Priede I.G. (2010)A large aggregation of liparids at 7703 meters and a reappraisal of abundance and diversity of hadal fishes. *BioScience* 60(7): 506-515.

Gianni, M. (2004) High seas bottom fisheries and their impact on the biodiversity of vulnerable deep-sea ecosystems: summary findings. IUCN, Gland, Switzerland: 83 pp.

Guinotte, J., Orr,J.,Cairns,S.,Freiwald,A.,Morgan,L.,George R.(2006).Will human-induced changes in seawater chemistry alter the distribution of deep-sea scleractinian corals? *Frontiers in Ecology and the Environment* 4(3): 141-146.

Harnik, P.G., Lotze, H.K.,Anderson,S.C.,

Liparid and macrourid fishes of the hadal zone: in situ observations of activity and feeding behaviour. *Proceedings of the Royal Society B: Biological Sciences* 276(1659): 1037-1045.

Jamieson, A.J., Lacey, N.L., Crz, A.N., Rowden, A.A., Piertney, S.B. (2013). The supergiant amphipod *Alicella gigantea* (Crustacea: Alicellida) from hadal depths in the Kermadec Trench, SW Pacific Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography* 92: 107-113.

Keeling R.F., Kortzinger A., Gruber, N. (2010). Ocean deoxygenation in a warming world. *Annual Review of Marine Science* 2: 199-229.

Koslow, J.A. (1996). Energetic and life history patterns of deep-sea benthic, benthopelagic and seamount-associated fishes. *Deep Sea Research* 43: 1081-1108.

Maier, C., Hegeman, W., Weinbauer, M.G., Gattuso J.P. (2009). Calcification of the cold-water coral *Lophelia pertusa*, under ambient and reduced pH. *Biogeosciences* 6: 1671-1680.

Matear, R.J., Hirst, A.C. (2003). Long-term changes in dissolved oxygen concentrations in the ocean caused by protracted global warming. *Global Biogeochemical Cycles* 17(4): 1125.

McClain, C.R., Lundsten, Ream, M., Barry, J., DeVogelaere, A. (2009). Endemicity, biogeography, composition, and community structure on a Northeast Pacific seamount. *PLOS Biology* 4 (1): e4141.

Menot, L., Sibuet, M., Carney, R.S., Levin, L.A., Rowe, G.T., Billett, D.S.M., Poore, G., Kitazato, H., Vanreusel, A., Galeron, J., Lavrado, H.P., Sellanes, J., Ingole, B.,

- Priede, I.G., Bergstad, O., Miller, P.I., Vecchione M., Gebruk, A., Falkenhaus T., Billett, D.S.M., Craig, J., Dale, A.C., Shields M.A. (2013). Does presence of a mid-ocean ridge enhance biomass and biodiversity? *PLoS ONE* 8(5): e61550.
- Ramirez-Llodra, E., Brand, A., Danovaro R., De Mol, B., Escobar E., German, C.R., Levin, L.A., Arbizu, P.M., Menot, L., Buhl-Mortensen, P., Narayanaswamy, B.E., Smith, C.R., Tittensor, D.P., Tyler, P.A., Vanreusel, A., Vecchione M. (2010). Deep, diverse and definitely different: unique attributes of the world's largest ecosystem *Biogeosciences* 7(9): 2851-2899.
- Ramirez-Llodra, E., Tyler, P.A., Baker, M.C., Bergstad, O.A., Clark, M.R., Escobar E., Levin, L.A., Menot, L., Rowden, A.A., Smith, C.R., Van Dover, C.L. (2011) Man and the last great wilderness: human impact on the deep sea. *PLoS ONE* 6(6): e22588.
- Richer de Forges, B., Koslow, J.A., Agre, G.C.B. (2000) Diversity and endemism of the benthic seamount fauna in the southwest Pacific. *Nature* 405: 944-947.
- Rogers, A.D. (1994) The biology of seamounts. *Advances in Marine Biology* 30: 305-350.
- Rogers, A.D., Baco, A., Griffiths, H., Hart, T., Hall-Spencer, J.M. (2007). Corals on seamounts. In: T.J. Pitcher, T. Morato, P.J.B. Hart et al. (Eds) *Seamounts:*

central Pacific: conservation challenges for future seabed mining. *Diversity and Distributions* DOI: 10.1111/ddi.12142.

Shaffer, G., Olsen, S., Medersea, O.P. (2009). Long-term ocean oxygen depletion in response to carbon dioxide emissions from fossil fuels. *Nature Geoscience* 105- 109.

Shepard, F.P. (1963). *Submarine Geology*. Harper & Row, New York, 557 pp.

Silva, H.M., Pinheiro, M.R. (2007) Small-scale fishing on seamounts. In: T.J. Pitcher, Morato, T., Hartet, P.J.B. et al (Eds). *Seamounts: Ecology, Fisheries & Conservation*. Blackwell Publishing, Oxford: 3360.

Smith, D.K. (1991) Seamount abundances and size distributions, and their geographic variations. *Reviews in Aquatic Sciences*

Marine Ecology Progress Series 442: 97

Troyanovsky, F.M., Lisovsky, S.F. (1995). Russian (USSR) fisheries research in deep waters (below 500 m) in the North Atlantic. In: A.G. Hopper (Ed). *Deep Water Fisheries of the North Atlantic Oceanic Slope*. Elsevier Academic Publishers, Dordrecht, Netherlands: 357-365.

Van Cauwenberghe, L., Vanreusel, M., Mees, J., Janssen, C.R. (2013). Microplastic pollution in deep-sea sediments. *Environmental Pollution* 182: 495-499.

Vecchione, M., Bergstad, O., Byrkjedal, J., Falkenhaus, T., Gebrek, A.V., Godø, O.R., Gislason, A., Heino, M., Høines, Å.S., Menezes, G.M., Piatkowski, U., Priede, I.G., Skov, H., Søiland, H., Sutton, T., Wenneck, T. (2010). Biodiversity patterns and processes on the mid-Atlantic Ridge. In: McIntyre, A.D. (Ed). *Life in the World's Oceans*. Blackwell Publishing Ltd: 361 pp.

Vetter, E.W., Smith, C.R., De Leo, F.C. (2010). Hawaiian hotspots: enhanced megafaunal abundance and diversity in submarine canyons on the oceanic islands of Hawaii. *Marine Ecology* 31(1): 183-199.

Vinogradova, N.G. (1997). Zoogeography of the abyssal and hadal zoobenthos. *Advances in Marine Biology* 32: 326-387.

Watson, R., Kitchingman, A., Cheung, W.W.L. (2007). Catches from world seamount fisheries. In: T.J. Pitcher, T. Morato, P.J.B. Hartet (Eds). *Seamounts: Ecology, Fisheries & Conservation*. Blackwell, Oxford: 400-412.

Webb, T.J., Beggs, E.V., O'Dor, R. (2010). Biodiversity's big wet secret: the global distribution of marine biological records reveals chronic underexploration of the deep pelagic ocean. *PLoS ONE*: e10223.

Wessel, P., Sandwell, D., Kim, S.S. (2010). The global seamount census. *Oceanography* 23: 24-33.

White, M., Bashmachnikov, A., Aristegui, J., Martins, A. (2007). Physical processes and seamount productivity. In: T.J. Pitcher, Morato, T., Hartet, P.J.B. (Eds). *Seamounts* 4(g)5(O2(.).3(c -053(ssTJ /TT12UI re)(u)-3 -1s(.).3(c -0v5(m)a -1.2240 Td)-3 O

23.51(ap)-1[(F72 Td [(W1.0021)79)14(ti)1UI4(larkl6c -0.001 Tw [(M)-4(o)-2(r20.00 T5T)-5(.,)-M-0.00R(1.0 -0.002 Tc

Wilson, R., Smith, K., Rosenblatt, R.H. (1985). Megafauna associated with bathyal seamounts in the central North Pacific Ocean. Deep Research 23: 1243-1254.

Wilson, R.R., Kaufman, R.S. (1987). Seamount biota and biogeography. In: B.H. Keating, P. Fryer, R. Batiza & G.W. Boehlert (