















the first time, the author has been able to find a single specimen of *Leptostomum* in the collections of the Royal Botanic Garden, Edinburgh.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is characterized by the presence of a single layer of trichomes, which are all of equal length, and by the fact that the leaves are all of the same size.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.

The genus *Leptostomum* is described by Schleicher in 1850, and the type species is *L. heterostichum* (L.) Schleicher.